

# The effect of desiccated soils on transpiration in spring wheat subject to elevated temperature and CO<sub>2</sub> concentration

Kobata Tohru

*Life and Environmental Science*

*Shimane University*

*Nisikawatu-cho, Matsue 690-8504, JAPAN*

## 1. INTRODUCTION

The dry matter production (DMP) for a time interval is indicated by

$$DMP = WUE \times T / VD \quad (1)$$

where WUE is water use efficiency, T transpiration and VD vapor deficit (Tanner and Sinclair, 1983). When plants are well irrigated, DMP<sub>0</sub>, WUE<sub>0</sub>, and T<sub>0</sub>/VD are given. The ratio of DMP to DMP<sub>0</sub> is

$$DMP/DMP_0 = (WUE/WUE_0) \times (T/VD) / (T_0/VD) \\ = (WUE/WUE_0) \times (T/T_0) \quad (2)$$

$$DMP = DMP_0 \times (WUE/WUE_0) \times (T/T_0) \quad (3)$$

In diverse crop plants it is expected that the WUE/WUE<sub>0</sub> scarcely changes with soil desiccation and is similar among cultivars but the T/T<sub>0</sub> is severely suppressed by soil desiccation (Tanner and Sinclair, 1983). Hence the response of T/T<sub>0</sub> to soil desiccation is one of the most important factors deciding the DMP. The DMP<sub>0</sub> is a potential productivity under well irrigated condition. The DMP<sub>0</sub> is expected to be estimated from a simulation model (Nakagawa and Horie 1997).

Empirically there was a close curve liner relationship between normalized transpiration rate (T/T<sub>0</sub>) and fraction of transpirable soil water (FTSW) in most crop species. The FTSW is ratio of soil water to transpirable soil water (Ray and Sinclair, 1998). Hence T/T<sub>0</sub> is the important factor to estimate the suppression of DMP<sub>0</sub> under soil desiccated conditions.

However, it is unknown the effect of soil water deficit on T/T<sub>0</sub> in spring wheat of Turkey subject to elevated temperature and CO<sub>2</sub> concentration.

## 2. MATERIALS and METHODS

### Plant materials

Adana99 of one of the dominant cultivar from Mediterranean area, Adana, Turkey was grown in pots

under glass house. Seedling soil for rice including fertilizer of 0.28 g of N, 0.33 g of P<sub>2</sub>O<sub>5</sub> and 0.3 g K<sub>2</sub>O per l (Green soil, Izumo Green Co.) was put into pots. Pot volume was 1.5 or 8 l. One or two plants were grown in a pot.

### Soil desiccation treatments

#### (1) Vegetative stage

In 2003 and 2004 plants were growth cabinet of Tottori Dryland Research Center after one week naturalization at the late vegetative stage from natural light glasshouse (Fig. 1- A&B). Temperature in ambient chamber was 20/15 °C, day time vapor deficit 11.9 g m<sup>-3</sup> and CO<sub>2</sub> concentration 414±20 μmol l<sup>-1</sup> and in an elevated chamber temperature 24/19 °C, day time vapor deficit 17.4 g m<sup>-3</sup> and CO<sub>2</sub> concentration 589±27 μmol l<sup>-1</sup>. Watering was stopped for two weeks.

In 2005 plants were grown in temperature and CO<sub>2</sub> controlled chambers. The chamber was covered with clear plastic film (Six-Light, Taiyo-Kogyo Co. Tokyo) and a temperature exchange fan and heater with a controller (Fig 1.-C). Temperature and CO<sub>2</sub> were controlled by several degrees higher than outside ambient and 600-700 μmol mol<sup>-1</sup>, respectively. Watering was terminated from the late vegetative stage for three weeks. Two chambers were used as replications.

#### (2) Reproductive stage

In 2006 plants grown in 8 l pot was used. Watering was withheld from the flowering stage to maturity in non temperature controlled glasshouse in ambient CO<sub>2</sub> concentration.

In both stages soil surface was covered with clear plastic beads to protect soil evaporation and pot weight was measured every day to estimate transpiration rate and soil water contents.

FTSW was calculated from the equation.

$$FTSW = SW_a / SW_{max} \quad (1)$$

where SW<sub>a</sub> is the actual water weight per pot and

SW<sub>0</sub>max is the available soil water weight per pot. SW<sub>max</sub> is decided by the difference between soil water weight at the field capacity and at the permanent wilting point. The field capacity is soil water weight at soil water potential of -0.03 MPa and the wilting point at -1.5 MPa (Kramer and Boyer, 1995). The soil moisture characteristic curve was decided for the soils with the thermocouple psychrometer methods (Boyer, 1995).


### 3. RESULTS and DISCUSSION

Transpiration rate decreased with a decrease of FTSW and there was a curve liner relationship between  $T/T_0$  and FTSW in both growth stage experiments (Fig. 1).

In the vegetative stage experiment the relationship under ambient and elevated conditions was similar within same experiment (Fig.1). However, precisely the curve should differ among experiments. Under climate change conditions plants suffer higher temperature and CO<sub>2</sub> concentrations from emergence and hence the plant may adapt to the environment. From the results the long term experiment in a chamber would be reliable because the plants have been elevated conditions since emergence (Fig. 1-C).

Actually wheat in the Mediterranean zone sometime suffers from terminal drought, that is, the soil desiccation during the grain-filling period (Turner, 1997). Drought response during the grain-filling period is the most important performance in wheat in Adana areas (personal communication by Dr. Genc). When watering was withheld from pots at flowering,  $T/T_0$  decreased with a decrease of FTSW similar to that at vegetative stage (Fig. 1-D). The result did not include the effect of elevated condition but the response of  $T/T_0$  to FTSW in the reproductive stage should not differ between ambient and elevated conditions due to that in the vegetative stage.

The results suggest that under elevated temperature and CO<sub>2</sub> conditions in the future the response of  $T/T_0$  to FTSW would be similar to that under ambient conditions and therefore the response curve should be available for sub-model to estimate the suppression of estimated biomass production by soil desiccation.



**Fig.1.** Transpiration ratio ( $T/T_0$ ) and FTSW in Adana99 under ambient (■) and elevated (□) temperature and CO<sub>2</sub> concentration during vegetative (A-C) and reproductive (D) stage. From A to C each point indicates average of three observations and for D an individual plant. The equation in D was used for a model.

### 4. REFERENCES

- Nakagawa, H., T. Horie, H.Y. Kim, H. Ohnishi and K. Homma (1997) Rice responses to elevated CO<sub>2</sub> concentrations and high temperature. *J. Agric. Meteorol.*52: 797-800.
- Ray, J. D. and Sinclair, T. R. 1998. *Ann. Bot.* 49:1381-1386.
- Turner, N. C. 1997. *Adv. Agron.* 58:293-338.
- Boyer, J. S. 1995. Measuring the water status of plants and soils. Academic Press, San Diego. USA.
- Kramer, P. J. and Boyer, J. S. 1995. Water relations of plants and soils. Academic Press, San Diego, USA.
- Tanner, C.B. and T.R. Sinclair 1983.1-28. In H.M.Taylor, et al. (ed) Limitations to efficient water use in crop production. Amer. Soc. Agron. Inc., Wisconsin.