

# Simulation of Crop Productivity for Evaluating Climate Change Effects

Tomohisa YANO<sup>1</sup>, Masumi KORIYAMA<sup>2</sup>, Tomokazu HARAGUCHI<sup>2</sup> and Mehmet AYDIN<sup>3</sup>

<sup>1</sup> Faculty of Engineering, Kyushu Kyouritsu University  
1-8 Jiyugaoka, Yahata-nishi-ku, Kitakyushu 807-8585, Japan

<sup>2</sup> Faculty of Agriculture, Saga University  
1 Honjo-cho, Saga 840-0027

<sup>3</sup> Faculty of Agriculture, Mustafa Kemal University  
31034 Antakya, Hatay, Turkey

## 1. Introduction

The effect of climate change on the crop productivity is usually investigated with the experimental methods using a growth chamber or with the numerical methods using a crop model.

The objectives of this research are (1) to establish the crop parameters in a crop growth model for wheat and maize as main crops in the world based on the experimental data collected in Adana and (2) to evaluate the effects of elevated CO<sub>2</sub> concentration and risen air temperature on crop growth.

## 2. Materials and methods

The SWAP model which was developed in the Netherlands (van Dam *et al.*, 1997) was used for the study. The model aims at simulating water, solute and heat transport in relation to plant growth at field scale level and for entire growing seasons. The SWAP model integrates soil-water balance and crop growth originally developed as the WOFOST model to describe daily phenological development and growth in response to environmental factors such as soils and climate, and crop management (Boogaard *et al.*, 1998). The model is eco-physiological process-based, simulating photosynthesis, evapotranspiration, and other major plant and soil processes. The major processes for crop growth are phenological development, CO<sub>2</sub>-assimilation, respiration, partitioning of assimilates to the various organs, and dry matter formation.

The SWAP requires input data on soil, crop and climate for its calibration and validation in the different environments. Climate (solar radiation, maximum and minimum temperatures, relative humidity, wind speed and rainfall), soil (soil water retention and hydraulic functions) and crop management data (crop calendar, some growth parameters, irrigation etc.) were collected for the

study location. The main parameters of the crop growth sub-model are temperature sum for the development rate of crop, specific leaf area, life span of leaves at optimum conditions, initial total crop dry weight, conversion of assimilates into biomass, fraction of dry matter increase partitioned to organs, maintenance respiration, reduction factor for senescence. Parameters for the crop growth model was decided with the crop growth measurement of two growing years for wheat and one growing year for maize.

The processes and relations incorporated in the WOFOST model are as follows: Using the absorbed radiation, which is absorbed by the canopy and is a function of incoming radiation and crop leaf area, and taking into account photosynthetic leaf characteristics the potential gross photosynthesis is calculated. The potential photosynthesis is reduced due to water and/or salinity stress, as quantified by the relative transpiration, and yields the actual gross photosynthesis. Part of the carbohydrates (CH<sub>2</sub>O) produced are used to provide energy for the maintenance of the existing live biomass (maintenance respiration). The remaining carbohydrates are converted into structural matter. In this conversion, some of the weight is lost as growth respiration. The dry matter produced is partitioned among roots, leaves, stems and storage organs, using partitioning factors that are a function of the phenological development stage of the crop. The fraction partitioned to the leaves, determines leaf area development and hence the dynamics of light interception. The dry weights of the plant organs are obtained by integrating their growth rates over time. During the development of the crop, part of living biomass dies due to senescence. Some simulated crop growth processes are influenced by temperature, like for example the maximum rate of photosynthesis and the maintenance respiration. Other processes, like the partitioning of assimilates or decay of crop tissue, are steered by the

phenological development stage (van Dam *et al.*, 1997).

The impacts of elevated CO<sub>2</sub> concentration and risen air temperature on crop growth and water balance components for wheat and second crop maize were examined using meteorological data of the Turkish Meteorological Station.


### 3. Results and discussions

Calculated and measured leaf area index (LAI) and biomass are compared in Fig. 1 for wheat in 2004/2005 and Fig. 2 for maize in 2004, respectively. The biomass values shown in these figures are total above-ground biomass. Although the discrepancy between calculated and measured values is recognized, estimation of crop growth with the SWAP model is satisfactory. The most sensitive parameters for crop growth simulation were found to be temperature sum for the development rate of crop, specific leaf area (i.e. the increase of the leaf area of the crop per kg weight increase of the living leaves) and initial crop total dry weight.


Since CO<sub>2</sub> is the fundamental parameter for crop growth prediction, it is supposed that use of only climatic data would cause the erroneous results.

Thus, effects of elevated CO<sub>2</sub> concentration on plants should be imposed to the model simulations. Maize is one of C<sub>4</sub> plants, and increases of CO<sub>2</sub> concentration are not so effective in crop growth unlike C<sub>3</sub> plants. Although wheat is a C<sub>3</sub> plant and elevated CO<sub>2</sub> concentration promotes photosynthesis, risen temperature may partially compensate it with stomatal closure. In fact, elevated atmospheric CO<sub>2</sub> concentration increases photosynthesis in C<sub>3</sub> plants and reduces evapotranspiration in both C<sub>3</sub> and C<sub>4</sub> plants due to reduced stomatal conductance.

Since CO<sub>2</sub> concentration is supposed to increase up to doubling concentration under SRES A2 scenario (IPCC 2001) in 2070s, simulation was done using 2004/2005 meteorological data under doubling CO<sub>2</sub> concentration condition. In doing calculation, the percent change in acclimatized photosynthesis rate was assumed to be +27% and +4% for wheat and maize, respectively (Cure and Acock, 1986). As the result, it was found that biomass values under the doubling CO<sub>2</sub> condition are about 17% and 5% higher than the values under the current CO<sub>2</sub> condition both for wheat and maize, respectively.



**Fig. 1.** Comparison between calculated LAI and biomass, and the measured ones of wheat.



**Fig. 2.** Comparison between calculated LAI and biomass, and the measured ones of maize.

However, high temperatures accelerate the phenology of plants, resulting in quicker maturation. The shortened growth cycle, in turn, may reduce the yield potential of annual crops (Rosenzweig and Hillel 1998). Based on a range of several current climate models, the mean annual global surface temperature is projected to increase by 1.4 to 5.8 °C over the period of 1990 to 2100 (IPCC, 2001). In order to evaluate the positive effect of elevated CO<sub>2</sub> concentration and the negative effect of risen air temperature on crop yields, simulation was done by doubling CO<sub>2</sub> concentration and by increasing maximum and minimum temperature by 1, 3, 5 °C relative to the current condition.

Results of calculated yields as well as evapotranspiration (ET) are shown in Table 1 and Table 2 for wheat and maize, respectively. Increase of biomass due to doubling CO<sub>2</sub> concentration is about 15% and 6% for wheat as a C3 crop and maize as a C4 crop, respectively. On the other hand, ET does not similarly increase and even decreases for maize, probably due to temporal water stress resulted from active crop growth. Decrease of biomass and ET following temperature rise is clearly shown in the both tables which denote that temperature increase of 3 °C results in biomass decreases by 17% and 20%, respectively. ET decreases similarly as biomass does.

Thus, increased yields by doubling CO<sub>2</sub> concentration are counteracted by rise in temperature of 3 and 1 °C, respectively for wheat and maize. More than 1 °C of temperature rise is estimated in 2070s when CO<sub>2</sub> concentration is supposed to increase up to doubling concentration, although much variations of temperature rise is recognized among models. Detailed description about temperature projection is given in the article “Prediction of Future Change of Water Demand Following Global Warming in the Cukurova Region,

Turkey” by Yano *et al.* of this issue.

Reduction of ET due to stomatal closure is not considered in Table 1 and 2. Experimental findings indicating considerable decreases in actual ET due to stomata closure under elevated CO<sub>2</sub> concentration have received a wide recognition (e.g. Ainsworth and Long, 2005). Decreases in transpiration by 17% and 26%, respectively for wheat and maize have been reported by Cure and Acock (1986). Further decreased ET values are replaced with those in Table 1 and 2 as shown in **Table 3**. Thus, ET from wheat and maize further decreases by 54 and 111 mm by transpiration reduction.

**Table 1** Effects of doubling CO<sub>2</sub> concentration (2xCO<sub>2</sub>) and rise in air temperature (+T) on crop growth and ET for wheat without ET reduction

Factor	Biomass (ton/ha)	Grain yield (ton/ha)	Growth days	ET (mm)
Control	17.74	4.71	164	370.4
2xCO <sub>2</sub>	21.67	5.74	162	371.4
+1 °C	16.41	4.14	155	350.3
+3 °C	13.21	3.96	141	310.4
+5 °C	10.37	4.01	128	254.3

**Table 2** Effects of doubling CO<sub>2</sub> concentration (2xCO<sub>2</sub>) and rise in air temperature (+T) on crop growth and ET for maize without ET reduction

Factor	Biomass (ton/ha)	Grain yield (ton/ha)	Growth days	ET (mm)
Control	27.14	17.29	120	378.4
2xCO <sub>2</sub>	28.85	17.98	120	359.0
+1 °C	25.13	15.83	116	361.7
+3 °C	21.71	13.14	110	335.2
+5 °C	18.62	10.79	107	314.4

#### 4. Conclusions

In this research, crop parameters in a detailed crop sub-model of the SWAP model were decided for wheat and maize using the field data of two growing years (2003/2004 and 2004/2005) for wheat and one growing year (2004) for maize in Adana. Consequently, the simulated values for LAI and biomass are in reasonable agreement with the measured data.

Then, the effects of elevated CO<sub>2</sub> concentration and risen air temperature on crop growth and evapotranspiration were separately evaluated using created meteorological data based on the observed data in 2004 and 2005.

Increase of biomass due to doubling CO<sub>2</sub> concentration is about 15% and 6% for wheat as a C3 crop and maize as a C4 crop, respectively. On the other hand, ET does not similarly increase and even decreases for maize. Decrease of biomass and ET following temperature rise is clearly shown that temperature increase of 3 °C results in biomass decrease for 17% and 20%, respectively. ET decreases similarly as biomass does.

Increased yields by doubling CO<sub>2</sub> concentration are counteracted by rise in temperature of 3 and 1 °C, respectively for wheat and maize. More than 1 °C of temperature rise is estimated in 2070s when CO<sub>2</sub> concentration is supposed to increase up to doubling concentration, although much variations of temperature rise is recognized among the different models.

#### Acknowledgements

We are grateful to Drs. M. Koc, M. Unlu and B. Bartcular of Cukurova University, Turkey for crop and meteorological data supply.

#### 5. Reference

- Ainsworth, E.A., Long, S.P., 2005. What have we learned from 15 years of free-air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *New Phytologist*, 165, 351-372.
- Boogaard, H. L., Van Diepen, C. A., Rötter, R. P., Cabrera, J. M. C. A., Van Laar, H. H., 1998. WOFOST 7.1. User's guide for the WOFOST 7.1 crop growth simulation model and WOFOST Control Center 1.5. Technical Document 52, DLO Winand Staring Centre, Wageningen, 142 pp.

**Table 3** Effects of ET reduction on crop growth and ET for wheat and maize at doubling CO<sub>2</sub> concentration (2xCO<sub>2</sub>)

Crop	Biomass (ton/ha)	Grain yield (ton/ha)	Growth days	ET (mm)
Wheat	22.06	6.15	162	317.8
Maize	28.44	17.74	120	248.2

- Cure, J. D., Acock, B., 1986. Crop responses to carbondioxide doubling: A literature survey. *Agric. For. Meteorol.*, 38: 127-145.
- IPCC, 2001. *Climate Change 2001: The Scientific Basis*. Cambridge University Press, Cambridge, 881pp.
- Rosenzweig, C., Hillel, D., 1998. *Climate Change and the Global Harvest*. Oxford University Press, New York, 324 pp.
- van Dam, J.C., Huygen, J., Wesseling, J. G., Feddes, R.A., Kabat, P., van Walsum, P. E. V., Groendijk, P., van Diepen, C. A., 1997. *Theory of SWAP version 2.0. Simulation of water flow, solute transport and plant growth in the Soil-Water-Atmosphere- Plant environment*. Technical Document 45, DLO Winand Staring Centre, Report 71, Dept. Water Resources, Agricultural University, Wageningen, 167 pp.